

## Propagating House Plants

Dale T. Lindgren, Department of Agronomy and Horticulture

The process of propagating various house plants is described in this NebGuide, which includes a listing of common house plants and how-to methods of propagating.

Propagating house plants in the home is an inexpensive and enjoyable hobby. Most house plants can be propagated at home satisfactorily with minimal special equipment.

House plants may be propagated by seeds, cuttings (tip and stem, cane, whole leaf, leaf section, leaf bud), division, runners, and air layering. All methods are described.

### Methods of Propagation

House plants may be propagated asexually, in which all new plants will be identical, in most cases, to the parent plant; or sexually by seed, where the new plants will not necessarily be identical to the parent plants. Cuttings, air layering, division, and runners are asexual methods of propagation.

**Seeds:** Some plants such as cactus and African violets can be raised in the home from seed. Plants grown from seeds take longer to reach maturity but are less costly than purchasing commercially available house plants. Variation among seedlings can make this method of propagation interesting.

There are many ways of germinating seed. Some are quite simple while others are complex. Basically, all are intended to provide seeds with the proper moisture and temperature conditions. Useful germinating media for seeds include sand, sphagnum moss, peat moss vermiculite, perlite, and a mixture of these media. The easiest method for germinating seed is to purchase commercial soil mixtures sold for germinating seed.

For germinating seed, follow the following steps:

1. Place the germinating medium in a container with a drainage hole in the bottom.
2. Moisten the medium by placing the container in a shallow pan of water. Remove when top of soil is moist.
3. Spread seeds in rows over surface of the medium.
4. Cover seeds according to directions on packet. Small seeds like African violets or begonia should not be covered.
5. Place the container in a polyethylene bag and seal the ends with tape or twist-ems.

6. Set the container on a window sill in indirect sun, with temperatures of 65° to 75°F.
7. Check the medium to maintain a damp condition.
8. Remove the bag when first seedling leaves are present. Provide maximum amount of light required by seedlings.
9. Transplant when first true leaves develop (usually the third leaf).

**Cuttings:** Cuttings are severed parent parts that produce roots and/or stems to form a new and independent plant. Stems, leaves, or roots may be used. Equipment needed for rooting cuttings include a container, rooting medium, a sharp knife, a plastic bag, a source of plant material, and in some cases a rooting hormone. A rooting hormone is useful for encouraging rooting on difficult-to-root cuttings.

Several methods of propagating plants by cuttings can be done. All are intended to provide the cuttings with the proper moisture and temperature conditions, just as with germinating seeds. Cuttings can be rooted in water, sand, perlite, or vermiculite. Some plants are easy to root in water, but perlite or vermiculite generally give more satisfactory results.

Select cuttings from healthy plants. When taking cuttings, make all cuts clean and at an angle through the stem, making sure there is at least one node (joint) under the surface of the medium. Push the cuttings down in the medium about 1 inch. The medium should be moist but not soggy. Slip an airtight polyethylene bag over the cuttings and around the container. No further watering will be necessary. Place in a room at 60° to 70°F. Cuttings can be potted when they show an abundance of roots.

### Types of Cuttings

**Tip and Stem Cuttings:** Tip cuttings (taken from the tip of plants) are used to propagate such common house plants as the velvet plant and jade plant. Tip cuttings generally are 3 to 5 inches long and are removed from the parent plant at a point just below a leaf (*Figure 1A*).

Swedish ivy, pilea, and fittonia can be propagated by stem cuttings (sections of stems with leaves attached). The cuttings should have three or four leaves for best rooting.

**Table I. List of common house plants and methods for propagating them. Some Species (spp) may be propagated by more than one method.**

African Violet ( <i>Saintpaulia</i> spp)	Seeds; whole leaf cuttings; division
Aloe ( <i>Aloe</i> spp)	Division
Arrowhead ( <i>Syngonium</i> spp)	Root cuttings that include a leaf node
Bloodleaf or Beefsteak Plant ( <i>Iresine</i> spp)	Tip cuttings
Boston Fern ( <i>Nephrolepis exaltata bostoniensis</i> )	Division; runners
Bromeliad (Many types)	Division
Cactus (Many types)	Seeds; whole leaf cuttings
Caladium ( <i>Caladium</i> spp)	Division
Christmas Cactus ( <i>Zygocactus</i> spp)	Tip cuttings
Croton ( <i>Codiaeum</i> spp)	Seeds; tip cuttings; air layering
Devil's Backbone ( <i>Pedilanthus tithymaloides variegatus</i> )	Stem cuttings
Dracaena, Red-margined ( <i>D. marginata</i> )	Stem cuttings; air layering; cane cuttings
Dracaena, Masange's ( <i>D. fragrans mauangeana</i> )	Stem cuttings; air layering; cane cuttings
Dracaena, Ribbon ( <i>D. sanderiana</i> )	Stem cuttings; air layering; cane cuttings
Dracaena, Gold Dust ( <i>JRGVHIQ</i> )	Stem cuttings; air layering; cane cuttings
Dumbcane ( <i>Dieffenbachia</i> spp)	Tip cuttings; cane cuttings; air layering; division
English Ivy ( <i>Hedera helix</i> )	Leaf bud cuttings
False Aralia ( <i>Dizygotheca</i> spp)	Seeds; air layering; stem cuttings
Fiddleleaf Fig ( <i>Ficus lyrata</i> )	Air layering
Fittonia ( <i>Fittonia</i> spp)	Stem cuttings
Jade Plant ( <i>Crassula</i> spp)	Tip cuttings; stem cuttings
Kalanchoe, Christmas ( <i>K. blossfeldiana</i> )	Leaf cuttings; stem cuttings
Kalanchoe, Dwarf Purple ( <i>K. pumila</i> )	Leaf cuttings; stem cuttings
Kalanchoe, Panda Plant ( <i>K. tomentosa</i> )	Leaf cuttings; stem cuttings
Kalanchoe <i>pinnata</i>	Plantlets on leaves
Kalanchoe <i>daigremontiana</i>	Plantlets on leaves
Kalanchoe <i>verticillata</i>	Plantlets on leaves
Moses in the Cradle ( <i>Rhoea</i> spp)	Division; seeds
Norfolk Island Pine ( <i>Araucaria</i> spp)	Seeds
Passion Flower ( <i>3DVVL</i> )	Seeds; stem cuttings
Peace Lily ( <i>Spathiphyllum wallisii</i> )	Division
Peperomia ( <i>Peperomia</i> spp)	Tip cuttings; whole leaf cuttings; leaf bud cuttings; division
Philodendron, Heartleaf ( <i>P. oxcardium</i> )	Stem cuttings
Philodendron, Velour ( <i>P. andreaum</i> )	Stem cuttings; air layering
Philodendron, Elephants Ear ( <i>P. domesticum</i> )	Stem cuttings; air layering
Philodendron, Fiddleleaf ( <i>P. panduraeformae</i> )	Stem cuttings; air layering
Philodendron, Splitleaf ( <i>P. pertusum</i> )	Stem cuttings; air layering
Pilea ( <i>Pilea</i> spp)	Stem cuttings; division
Poinsettia ( <i>Euphorbia pulcherrima</i> )	Tip cuttings (allow wound to heal)
Prayer Plant ( <i>Marantha</i> spp)	Division
Rubber Plant ( <i>Ficus elastica</i> )	Air layering; seeds
Schefflera ( <i>Brassaia</i> spp)	Seeds; cuttings of half ripened stems
Sentry Palm ( <i>Howeia</i> spp)	Seeds
Snake Plant ( <i>Sansevieria</i> spp)	Leaf section cuttings; division
Spider Plant ( <i>Chlorophytum</i> spp)	Division; runners
Swedish Ivy ( <i>Plectranthus</i> spp)	Stem cuttings
Umbrella Plant ( <i>Cyperus</i> spp)	Seeds; division; upside down stem cuttings
Velvet Plant ( <i>Gynura</i> spp)	Tip cuttings; stem cuttings




Figure 1. Tip and stem cuttings (A), and cane cuttings (B).

**Cane Cuttings:** Cane cuttings are used for propagating dumbcane, Chinese evergreen, and similar plants that produce cane-like or leafless stems. The cane is cut up into small pieces 2 to 3 inches long. Place the cuttings on their sides slightly below the surface of the rooting medium. A bud eventually will sprout and form a new stem when the cutting is rooted (Figure 1B).

**Whole Leaf Cuttings:** Whole leaf cuttings are prepared from leaves with or without their stalks (called petioles). Roots and leaves eventually will form at the base of the leaf (Figure 2A). Peperomia and African violets commonly are started by whole leaf cuttings.




Figure 2. Whole leaf cuttings (A) and leaf section cuttings (B).

**Leaf Section Cuttings:** Leaf section cuttings can be used for propagating plants like the Rex begonia and snake plant. The leaves are cut into pieces, with the edge of the cuttings closest to the base of the parent plant inserted into the rooting medium (Figure 2B).

**Leaf Bud Cuttings:** Leaf bud cuttings consist of a single leaf attached to a piece of 1- to 1½-inch stem. The dormant bud, located where the leaf stalk joins the stem will give rise to a new shoot and branches (Figure 3). The cutting should be inserted in the rooting medium with the bud about ½ inch below the surface. English ivy easily is propagated by this method.




Figure 3. Leaf bud cuttings.

Cuttings from succulents or cactus should be allowed to dry for 1 to 7 days, depending upon species and size, before placing in a rooting medium. The drying period will cause the cut edges to callous. This will prevent the absorption of excessive amounts of moisture that could result in rotting.

### Division

Division is the quickest way to propagate plants that produce stems at their base. To divide a plant, remove it from the container and separate the plant into smaller pieces with a knife or fingers. Each division should have some roots. Divisions without roots may be rooted as tip cuttings. Plants can be divided at any time, although the dormant or rest period is best. Use division rather than leaf cuttings to propagate variegated plants, such as sansevieria. The non-variegated type is produced from leaf cuttings.

### Runners

Runners, which are prostrate creeping shoots or stems such as found on strawberry geraniums, can be used to propagate some house plants. Place the leafy cluster of the runners in contact with a sandy loam soil in a second pot and hold them




Figure 4. Propagating plant runners.

down with a paperclip or similar item (Figure 4). When roots have developed from the cluster, the new plant can be cut free from the runner. A simpler method would be to cut off the leafy clusters along with a small section of stem and root them as tip cuttings.

### Air layering

Air layering is a method extremely useful in shortening certain plants which become “leggy.” With a sharp knife, make an upward or downward slanting cut 1- to 1½-inches long and about one-quarter to one-half way through the stem. Insert a matchstick or toothpick into the wound to prevent it from healing over and dust a small amount of rooting hormone, if available, into the incision (Figure 5A). The rooting hormone usually hastens rooting if applied correctly, although some plants root just as well untreated. Remove leaves or twigs on




Figure 5. Put a matchstick or toothpick into the cut stem when air layering (A), and wrap with sphagnum moss (B).

the chosen stem about 3 to 4 inches above the incision point. Wrap the wound with two handfuls of moist, but not soggy, sphagnum moss to form a ball around the stem (Figure 5B). Wrap the moss with plastic film (example, a freezing bag) and seal the seam and edges with cellophane or electrical plastic waterproof tape or twist-ems to prevent water loss from the moss (Figure 6A).

After several weeks, roots should be visible near the edges of the moss ball under the plastic sheet. When a good root system has formed, remove the plastic film and sever the rooted tip from the stock plant at a point just below the ball of moss and roots (Figure 6B). Pot the rooted layer in the recommended soil mixture. The remaining portion of the cane can be saved and new buds will arise on it.

These are the most common methods of propagating house plants in the home. Many Internet sources are available detailing methods of propagation for specific house plants.




Figure 6. Wrap sphagnum moss with plastic and seal edges (A), and sever rooted tip from plant stock when root system is developed (B).

### Acknowledgment

The author would like to recognize contributions made by Don Steinegger, retired UNL horticulturist, to the first edition of this publication.

**This publication has been peer reviewed.**

UNL Extension publications are available online at <http://extension.unl.edu/publications>.

**Index: Lawn & Garden  
Ornamentals**

Issued May 2008

Extension is a Division of the Institute of Agriculture and Natural Resources at the University of Nebraska–Lincoln cooperating with the Counties and the United States Department of Agriculture.

University of Nebraska–Lincoln Extension educational programs abide with the nondiscrimination policies of the University of Nebraska–Lincoln and the United States Department of Agriculture.

© 2008, The Board of Regents of the University of Nebraska on behalf of the University of Nebraska–Lincoln Extension. All rights reserved.